

UNKNOTTING INFORMATION FROM HEEGAARD FLOER HOMOLOGY

BRENDAN OWENS

ABSTRACT. We use Heegaard Floer homology to obtain bounds on unknotting numbers. This is a generalisation of Ozsváth and Szabó's obstruction to unknotting number one. We determine the unknotting numbers of 9_{10} , 9_{13} , 9_{35} , 9_{38} , 10_{53} , 10_{101} and 10_{120} ; this completes the table of unknotting numbers for prime knots with crossing number nine or less. Our obstruction uses a refined version of Montesinos' theorem which gives a Dehn surgery description of the branched double cover of a knot.

1. INTRODUCTION

Let K be a knot in S^3 . Given any diagram D for K , a new knot may be obtained by changing one or more crossings of D . The unknotting number $u(K)$ is the minimum number of crossing changes required to obtain the unknot, where the minimum is taken over all diagrams for K .

Let $\Sigma(K)$ denote the double cover of S^3 branched along K . A theorem of Montesinos ([9], or see Lemma 3.1) tells us that for any knot K , $\Sigma(K)$ is given by Dehn surgery on some framed link in S^3 with $u(K)$ components, with half-integral framing coefficients. In particular if $u(K) = 1$ then $\Sigma(K)$ is obtained by $\pm \det K/2$ Dehn surgery on a knot C , where $\det K$ is the determinant of K . Ozsváth and Szabó have shown in [16] that the Heegaard Floer homology of a 3-manifold Y gives an obstruction to Y being given by half-integral surgery on a knot in S^3 ; they apply this to $\Sigma(K)$ to obtain an obstruction to K having unknotting number one.

Note that crossings in a knot diagram may be given a sign as in Figure 1 (independent of the choice of orientation of the knot). Let $\sigma(K)$ denote the signature of a knot K . It is shown in [3, Proposition 2.1] (also [17, Theorem 5.1]) that if K' is obtained from K by changing a positive crossing, then

$$\sigma(K') \in \{\sigma(K), \sigma(K) + 2\};$$

similarly if K' is obtained from K by changing a negative crossing then

$$\sigma(K') \in \{\sigma(K), \sigma(K) - 2\}.$$

Now suppose that K may be unknotted by changing p positive and n negative crossings (in some diagram). Since the unknot has zero signature, it follows that a bound




FIGURE 1. Signed crossings in a knot diagram.

for n is given by

$$(1) \quad n \geq \sigma(K)/2.$$

The main result of this paper is an obstruction to equality in (1). This is easiest to state for the case of an alternating knot; the obstruction is then a condition on the Goeritz matrix obtained from an alternating projection of K . (We will recall the definition of the Goeritz matrix in Section 4.) We also restrict for now to knots which can be unknotted with two crossing changes.

A positive-definite integral matrix Q of rank r presents a finite group Γ_Q via the short exact sequence

$$0 \longrightarrow \mathbb{Z}^r \xrightarrow{Q} \mathbb{Z}^r \longrightarrow \Gamma_Q \longrightarrow 0.$$

A *characteristic covector* for Q is an element of $\mathbb{Z}^r$ which is congruent modulo 2 to the diagonal of Q , i.e., an element of

$$\text{Char}(Q) = \{\xi \in \mathbb{Z}^r \mid \xi_i \equiv Q_{ii} \pmod{2}\}.$$

Suppose that $\det Q$ is odd. Define a function

$$m_Q : \Gamma_Q \rightarrow \mathbb{Q}$$

by

$$m_Q(g) = \min \left\{ \frac{\xi^T Q^{-1} \xi - r}{4} \mid \xi \in \text{Char}(Q), [\xi] = g \right\}.$$

(The minimum exists since Q is positive-definite.)

Theorem 1. *Let K be an alternating knot which may be unknotted by changing p positive and n negative crossings, where $n = \sigma(K)/2$ and $p + n = 2$. Let G be the positive-definite Goeritz matrix obtained from an alternating diagram for K . Then*

there exists a positive-definite matrix

$$\tilde{Q} = \begin{pmatrix} m_1 & 1 & a & 0 \\ 1 & 2 & 0 & 0 \\ a & 0 & m_2 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix},$$

with

$$\det \tilde{Q} = \det K,$$

$$0 \leq a < m_1 \leq m_2 \quad (\text{and hence } a < \det K/4),$$

and exactly n of $\{m_1, m_2\}$ are even; and a group isomorphism

$$\phi : \Gamma_{\tilde{Q}} \rightarrow \Gamma_G$$

with

$$m_{\tilde{Q}}(g) \geq m_G(g),$$

$$\text{and } m_{\tilde{Q}}(g) \equiv m_G(g) \pmod{2}$$

for all $g \in \Gamma_{\tilde{Q}}$.

Applying Theorem 1 to the alternating knots which were listed in [1] as having unknotting number 2 or 3 yields the following:

Corollary 2. *The knots $9_{10}, 9_{13}, 9_{35}, 9_{38}, 10_{53}, 10_{101}, 10_{120}$ have unknotting number 3.*

For all but one of the knots in Corollary 2, the signature is 4 and the unknotting number computation follows directly from Theorem 1. The exception is 9_{35} , whose signature is 2. The computation of $u(9_{35})$ uses Theorem 1 and also a result of Traczyk [18].

Corollary 2 completes the table of unknotting numbers for prime knots with 9 crossings or less.

Recall that for an oriented framed link $C_1, \dots, C_r$ in S^3 , the linking matrix is the symmetric matrix (a_{ij}) with each diagonal entry a_{ii} given by the framing on C_i , and off-diagonal entries a_{ij} given by the linking numbers $\text{lk}(C_i, C_j)$. The following is a refinement of Montesinos' theorem which was inspired by a theorem of Cochran and Lickorish [3, Theorem 3.7].

Theorem 3. *Suppose that a knot K may be unknotted by changing p positive and n negative crossings, with $n = \sigma(K)/2$. Then the branched double cover $\Sigma(K)$ may be obtained by Dehn surgery on an oriented, framed $p + n$ component link $C_1, \dots, C_{p+n}$ in S^3 with linking matrix $\frac{1}{2}Q$, where Q is a positive-definite integral matrix which is congruent to the identity modulo 2, and exactly n of the diagonal entries of Q are congruent to 3 modulo 4.*

Moreover, by handlesliding and changing orientations one may replace the linking matrix with $\frac{1}{2}PQP^T$, for any $P \in GL(p + n, \mathbb{Z})$ which is congruent to the identity modulo 2. This preserves the congruences modulo 4 on the diagonal.

It is shown in [12] that the double branched cover of the Montesinos knot 10_{145} does not bound any positive-definite four-manifold. This knot has signature two. Combining this with Theorem 3 (or the above-mentioned theorem of Cochran and Lickorish) yields the following:

Corollary 4. *If 10_{145} is unknotted by changing p positive crossings and n negative crossings, then $n \geq 2$.*

Given a matrix Q in $M(r, \mathbb{Z})$ which is conjugate modulo 2 to the identity, associate a matrix $\tilde{Q} \in M(2r, \mathbb{Z})$ by replacing each entry by a 2×2 -block as follows:

$$(2) \quad \begin{aligned} \text{odd entries:} \quad 2m-1 &\mapsto \begin{bmatrix} m & 1 \\ 1 & 2 \end{bmatrix} \\ \text{even entries:} \quad 2a &\mapsto \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Thus for example if $r = 2$,

$$Q = \begin{pmatrix} 2m_1-1 & 2a \\ 2a & 2m_2-1 \end{pmatrix} \mapsto \tilde{Q} = \begin{pmatrix} m_1 & 1 & a & 0 \\ 1 & 2 & 0 & 0 \\ a & 0 & m_2 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix}.$$

For a rational homology three-sphere Y , the *correction terms* of Ozsváth and Szabó are a set of rational numbers $\{d(Y, \mathfrak{s}) \mid \mathfrak{s} \in \text{Spin}^c(Y)\}$ which provide constraints on which four-manifolds Y may bound. We recall these constraints in Section 4; combining these with Theorem 3 yields the following unknotting obstruction, of which Theorem 1 is a special case.

Theorem 5. *Let K be a knot in S^3 which may be unknotted by changing p positive and n negative crossings, where $n = \sigma(K)/2$. Let $Q_1, \dots, Q_k$ be a complete set of representatives of the finite quotient*

$$\frac{\{Q \in M(p+n, \mathbb{Z}) \mid Q \text{ is positive-definite, } \det Q = \det K, Q \equiv I \pmod{2}\}}{\{P \in GL(p+n, \mathbb{Z}) \mid P \equiv I \pmod{2}\}},$$

and let $\tilde{Q}_1, \dots, \tilde{Q}_k$ be the corresponding elements of $M(2(p+n), \mathbb{Z})$. Then for some Q_i which has exactly n diagonal entries conjugate to 3 modulo 4, there exists a group isomorphism

$$\phi : \Gamma_{\tilde{Q}_i} \rightarrow \text{Spin}^c(\Sigma(K))$$

with

$$\begin{aligned} m_{\tilde{Q}_i}(g) &\geq d(\Sigma(K), \phi(g)), \\ \text{and } m_{\tilde{Q}_i}(g) &\equiv d(\Sigma(K), \phi(g)) \pmod{2} \end{aligned}$$

for all $g \in \Gamma_{\tilde{Q}_i}$.

The following example illustrates the use of Theorem 5 to obstruct higher unknotting numbers.

Corollary 6. *The 11-crossing two-bridge knot $S(51, 35)$ has unknotting number 4.*

Acknowledgements. The problem of generalising the obstruction in [16] to higher unknotting numbers was suggested to me by Peter Ozsváth. I am grateful to Peter Ozsváth, Ravi Ramakrishna, and Sašo Strle for helpful discussions. Some Maple programs used in verifying Corollaries 2 and 6 were written jointly with Sašo Strle.

2. KIRBY-ROLFSEN CALCULUS

In this section we establish some preliminaries on Dehn surgery. For details on Dehn surgery and Kirby-Rolfsen calculus see [5].

A framed link L in S^3 with rational framing coefficients determines a three-manifold Y_L by Dehn surgery (remove a tubular neighbourhood of each component of L ; the framing coefficient determines the gluing map to sew back a solid torus along the boundary). If the framing coefficients are integers one obtains a four-manifold W_L with boundary Y_L by attaching two-handles to B^4 along the components of L . Kirby-Rolfsen calculus describes when two framed links L, L' determine the same three-manifold Y_L .

Given a framed oriented link L with components $C_1, \dots, C_m$, let A denote the free abelian group with generators $c_1, \dots, c_m$. Define a symmetric bilinear form

$$Q : A \times A \rightarrow \mathbb{Q}$$

by

$$Q(c_i, c_j) = \begin{cases} \text{framing coefficient of } C_i & \text{if } i = j; \\ \text{linking number } \text{lk}(C_i, C_j) & \text{if } i \neq j. \end{cases}$$

In other words, the matrix of Q in the basis $c_1, \dots, c_m$ is the linking matrix of L . (This is the intersection pairing on $H_2(W_L; \mathbb{Z})$ if the diagonal entries are integers.)

In the case that the framing coefficients on L are integers, any change of basis in A may be realised by a change in the link L . In particular the change of basis $c_i \mapsto c_i \pm c_j$ may be realised by a *handleslide*. Let λ_j denote a pushoff of C_j whose linking number with C_j equals the framing of C_j . A handleslide $C_i \mapsto C_i \pm C_j$ consists of replacing C_i by the oriented band sum of C_i with $\pm \lambda_j$. This gives a new link L' whose linking matrix is the matrix of Q in the basis $c_1, \dots, c'_i = c_i \pm c_j, \dots, c_m$ and with $Y_{L'} \cong Y_L$, $W_{L'} \cong W_L$. It will be convenient to have the following generalisation of handlesliding to links with rational framings.

Proposition 2.1. *Let L be an oriented link in S^3 consisting of components $C_1, \dots, C_m$ with framings $\frac{p_1}{q_1}, \dots, \frac{p_m}{q_m}$, and let Q be the rational-valued bilinear pairing determined by the linking matrix of L . Then by replacing C_i in L it is possible to obtain a link L' whose linking matrix is the matrix of Q in the basis $c_1, \dots, c'_i = c_i \pm q_j c_j, \dots, c_m$ and with $Y_{L'} \cong Y_L$.*

Proof. For each $j = 1, \dots, m$ choose a continued fraction expansion

$$\frac{p_j}{q_j} = a_{l_j}^j - \frac{1}{a_{l_j-1}^j - \dots - \frac{1}{a_1^j}}.$$

(The numbers $a_{l_j}^j, \dots, a_1^j$ arise from the Euclidean algorithm as follows:

$$\begin{aligned} r_{l_j} = p_j &= a_{l_j}^j q_j - r_{l_j-2} \\ r_{l_j-1} = q_j &= a_{l_j-1}^j r_{l_j-2} - r_{l_j-3} \\ &\vdots \\ r_2 &= a_2^j r_1 - 1 \\ r_1 &= a_1^j. \end{aligned} \tag{3}$$

Use reverse “slam-dunks” to obtain an integral surgery description of Y_L : as shown in Figure 2, we add a chain of linked unknots linking each C_j , with framings $a_1^j, \dots, a_{l_j-1}^j$, and replace the framing on C_j with $a_{l_j}^j$. (This is a standard procedure, see e.g. [5, §5.3].) Denote the resulting link by $L_{\mathbb{Z}}$, and let $Q_{\mathbb{Z}} : A_{\mathbb{Z}} \times A_{\mathbb{Z}} \rightarrow \mathbb{Z}$ denote the resulting bilinear form.




FIGURE 2. Converting Dehn surgery to integral surgery.

We now perform handleslides on this integer-framed link. Let $U_1, \dots, U_{l_j-1}$ be the chain of unknots linking C_j as above, oriented so that $\text{lk}(C_j, U_{l_j-1}) = \text{lk}(U_k, U_{k-1}) = -1$, for $2 \leq k < l_j$. Let $K_1 = C_i + U_1$, and note that

$$\begin{aligned} \text{lk}(K_1, U_1) &= a_1^j, \\ \text{lk}(K_1, U_2) &= -1. \end{aligned} \tag{4}$$

We now define K_k recursively for $2 \leq k < l_j$. Choose any link diagram of $K_{k-1} \cup U_{k-1} \cup U_k$. By performing a handleslide over U_k for each crossing where K_{k-1} crosses over U_{k-1} we obtain a knot K_k which does not cross over U_{k-1} and therefore is separated from it by a two-sphere in S^3 (see Figure 3). The signed count of these handleslides is equal to the linking number of K_{k-1} and U_{k-1} ; thus we write

$$[K_k] = [K_{k-1}] + \text{lk}(K_{k-1}, U_{k-1})[U_k],$$

where $[K_k]$ denotes the element of $A_{\mathbb{Z}}$ corresponding to the knot K_k . We may use this to compute linking numbers and the framing of K_k . In particular

$$(5) \quad \text{lk}(K_2, U_2) = -1 + a_2^j \text{lk}(K_1, U_1),$$

and for $2 < k < l_j$,

$$(6) \quad \begin{aligned} \text{lk}(K_k, U_k) &= \text{lk}(K_{k-1}, U_k) + a_k^j \text{lk}(K_{k-1}, U_{k-1}) \\ &= -\text{lk}(K_{k-2}, U_{k-2}) + a_k^j \text{lk}(K_{k-1}, U_{k-1}). \end{aligned}$$

Finally we let C'_i be obtained as above from K_{l_j-1} by sliding over C_j , with C'_i unlinked from each of $U_1, \dots, U_{l_j-1}$. We then have

$$(7) \quad \begin{aligned} [C'_i] &= [K_{l_j-1}] + \text{lk}(K_{l_j-1}, U_{l_j-1})[C_j], \\ \text{lk}(C'_i, C_j) &= -\text{lk}(K_{l_j-2}, U_{l_j-2}) + a_{l_j}^j \text{lk}(K_{l_j-1}, U_{l_j-1}) + \text{lk}(C_i, C_j). \end{aligned}$$

Comparing (4), (5), (6), and (7) to (3) we see that

$$\begin{aligned} \text{lk}(K_k, U_k) &= r_k^j \quad \text{for } k = 1, \dots, l_j - 2, \\ \text{lk}(K_{l_j-1}, U_{l_j-1}) &= r_{l_j-1}^j = q_j, \\ \text{lk}(C'_i, C_j) &= p_j + \text{lk}(C_i, C_j). \end{aligned}$$

This yields

$$[C'_i] = [C_i] + \mathcal{U} + q_j[C_j],$$

where

$$\mathcal{U} = [U_1] + \sum_{k=2}^{l_j-1} r_{k-1}^j [U_k].$$

Note that by construction C'_i is separated by a two-sphere from each U_k and so $Q_{\mathbb{Z}}([C'_i], \mathcal{U}) = 0$. The framing of C'_i is given by

$$\begin{aligned} Q_{\mathbb{Z}}([C'_i], [C'_i]) &= Q_{\mathbb{Z}}([C_i] + \mathcal{U} + q_j[C_j], [C_i] + \mathcal{U} + q_j[C_j]) \\ &= Q_{\mathbb{Z}}([C_i] + q_j[C_j], [C_i] + \mathcal{U} + q_j[C_j]) \\ &= Q_{\mathbb{Z}}([C_i], [C_i]) + 2q_j Q_{\mathbb{Z}}([C_i], [C_j]) + q_j^2 a_{l_j}^j - q_j r_{l_j-2} \\ &= a_{l_i}^j + 2q_j \text{lk}(C_i, C_j) + p_j q_j. \end{aligned}$$




FIGURE 3. **Handlesliding K_{k-1} over U_k yields K_k which is separated from U_{k-1} by a two-sphere.**

Slam dunking to remove the chains of linking unknots from each of $C_1, \dots, C'_i, \dots, C_m$ gives the required link L' for the basis change $c'_i = c_i + q_j c_j$. To get the opposite sign construct C'_i as above but start with $K_1 = C_i - U_1$. $\square$

The following lemma is an application of the standard procedure, referred to in the proof of Proposition 2.1 and illustrated in Figure 2, for converting a Dehn surgery description of a three-manifold to an integral surgery description.

Lemma 2.2. *Let $L = \{C_1, \dots, C_r\}$ be a framed link in S^3 with framing $(2m_i - 1)/2$ on C_i , and let Y be the three-manifold obtained by Dehn surgery on L . Then Y is equal to the boundary of the four-manifold W obtained by adding 2-handles to B^4 along either of the following $2n$ -component framed links (as in Figure 4):*

- (i) *the link consisting of the components C_i with framing m_i plus a small linking unknot with framing 2, for each $i = 1, \dots, r$;*
- (ii) *the link consisting of C_i with framing m_i , plus a longitude C'_i with framing m_i and with the opposite orientation, with linking number $\text{lk}(C_i, C'_i) = 1 - m_i$, for each $i = 1, \dots, r$.*

Proof. The fact that Y is the boundary of the four-manifold given by the framed link (i) follows from the continued fraction expansions $(2m_i - 1)/2 = m_i - \frac{1}{2}$. The

equivalence between (i) and (ii) follows by handlesliding: add C_i to C'_i to go from (ii) to (i). $\square$

Recall that to each matrix $Q \in M(r, \mathbb{Z})$ which is congruent to the identity modulo 2, we associate the matrix $\tilde{Q} \in M(2r, \mathbb{Z})$ as in (2). If a 3-manifold Y is given by Dehn surgery on a link with linking matrix $\frac{1}{2}Q$, then by Lemma 2.2, Y is the boundary of a simply-connected four-manifold with intersection pairing $\tilde{Q}$. Also note that $\det Q = \det \tilde{Q}$, and Q is positive-definite if and only if $\tilde{Q}$ is positive-definite: let

$$\Delta_k(Q) = \det(Q_{ij})_{i,j \leq k}.$$

Then

$$\begin{aligned} \Delta_{2k}(\tilde{Q}) &= \Delta_k(Q), \\ \Delta_{2k-1}(\tilde{Q}) &= (\Delta_{2k-2}(\tilde{Q}) + \Delta_{2k}(\tilde{Q}))/2. \end{aligned}$$




FIGURE 4. **Half-integer surgery.** There are $2m_i - 2$ crossings in the diagram on the right.

3. PROOF OF THEOREM 3

The proof of Theorem 3 consists of three lemmas. The first of these is a proof of Montesinos' theorem using Kirby calculus. We could omit this and simply refer to proofs in the literature, for example [7] (or to the proof of Lemma 3.2). We include the proof since the four-dimensional point of view initially led us to a proof of Theorem 3, and since it spells out a useful algorithm for drawing a surgery diagram of $\Sigma(K)$. (For more details on Kirby diagrams of cyclic branched covers see [5, §6.3]; indeed what follows is a variation of the method in their Exercise 6.3.5(c).)

Lemma 3.1. *Let K be a knot in S^3 which can be unknotted by changing r crossings in some diagram D . Then the double branched cover $\Sigma(K)$ is given by Dehn surgery on an r -component link in S^3 with linking matrix $\frac{1}{2}Q$, where Q is congruent to the identity modulo 2.*

Proof. We think of $K \subset S^3$ as being in the boundary of B^4 . Draw r unlinked unknots beside D , each with framing $+1$. This is a Kirby diagram which represents K as a knot in the boundary of the “blown up” four-ball $X = B^4 \# r\mathbb{CP}^2$. As observed in [3], the knot K bounds a disk Δ in X . This may be seen from the diagram by sliding each of the chosen crossings in D over a $+1$ -framed unknot as in Figure 5. Mark each of these changed crossings with a small arc α_i , $i = 1, \dots, r$, as shown in that figure.




FIGURE 5. Changing crossings by sliding over a two-handle.

The resulting diagram consists of:

- an unknot U which has been obtained from K by crossing changes;
- arcs $\alpha_1, \dots, \alpha_r$ (one per changed crossing);
- $+1$ -framed unknots $\gamma_1, \dots, \gamma_r$.

Each γ_i bounds a disk D_i which retracts onto α_i and whose intersection with U consists of the endpoints of α_i .

It is also observed in [3] that $H_1(X - \Delta; \mathbb{Z}/2) \cong \mathbb{Z}/2$, with generator given by the meridian of K . (To see this note from Figure 5 that the linking number of U with each of the $+1$ -framed unknots is even. Now use the Mayer-Vietoris sequence for the decomposition of X into $X - \Delta$ and a neighbourhood of Δ , with $\mathbb{Z}/2$ coefficients.) Thus there exists a unique double cover W of X branched along Δ ; this is a four-manifold with boundary $\Sigma(K)$.

Rearrange the diagram so that a point of U which is not the endpoint of an arc α_i is the point at infinity and U is a vertical line; then Δ may be seen in this diagram as the half-plane to the left of U . (For a simple example see the first 3 diagrams in




FIGURE 6. **Drawing the double branched cover.** Here Δ is the half-plane to the left of U . The dotted line in the diagrams on the right is the preimage of U . The top two diagrams occur at endpoints of α_i , and the bottom two occur where α_i intersects the interior of Δ .

Figure 7. Note in general the arcs α_i may be knotted and linked, and may intersect Δ .) We may rearrange the diagram so that all intersections of γ_i and Δ look like one of the diagrams on the left of Figure 6. To draw a Kirby diagram of W , we simply need to take two copies of $S^3 - U$ cut open along Δ , and join the boundary half-planes in pairs. Or in other words: take the part of the diagram to the right of U , and draw another copy of it to the left of U . (Think of rotating the half plane to the right of U about U by π , *not* reflecting.) Complete the centre of the diagram using Figure 6. (For an example see Figure 7.)




FIGURE 7. A four-manifold bounded by the double branched cover of the left-handed trefoil.

Each arc α_i lifts to a knot $\tilde{\alpha}_i$, and each D_i lifts to an annulus $\tilde{D}_i$ with core $\tilde{\alpha}_i$. The knot γ_i lifts to two knots C_i, C'_i ; these are the boundary of the annulus $\tilde{D}_i$.

We now compute the framings of C_i, C'_i . The 0-framing on γ_i lies on the disk D_i , and lifts to a curve on the annulus $\tilde{D}_i$. This is the same framing for C_i (or C'_i) as the other boundary curve of $\tilde{D}_i$, but with the opposite sign. Thus the 0-framing on γ_i lifts to the $-\text{lk}(C_i, C'_i)$ -framing on each of C_i, C'_i . Then the framing $+1$ on γ_i lifts to m_i on each of C_i, C'_i , where $\text{lk}(C_i, C'_i) = 1 - m_i$.

We note that the resulting Kirby diagram for W matches that in Lemma 2.2 (ii). That lemma then shows that $\Sigma(K) = \partial W$ is Dehn surgery on the framed link $L = C_1, \dots, C_r$ with framing $(2m_i - 1)/2$ on C_i . $\square$

To prove that the matrix Q is positive-definite under the hypotheses of Theorem 3 one may appeal to [3, Theorem 3.7], which gives a formula for the signature of the four-manifold W constructed in Lemma 3.1. Surprisingly however it is also possible to prove this using the following purely three-dimensional argument.

Lemma 3.2. *Suppose that K may be unknotted by changing p positive and n negative crossings, with $n = \sigma(K)/2$. Let $\Sigma(K)$ be given by Dehn surgery on a link $C_1, \dots, C_{p+n}$ with linking matrix $\frac{1}{2}Q$ as in Lemma 3.1. Then Q is positive-definite, and exactly n of the diagonal entries of Q are congruent to 3 modulo 4.*

Proof. The positivity of Q is proved in [16, Theorem 8.1] for the case of unknotting number one knots, i.e. $p + n = 1$. We include the proof of this case here for completeness.




FIGURE 8.

Suppose K_- , K_0 and K_+ are links in S^3 which are identical outside of a ball in which they appear as in Figure 8. Recall that the double cover of a ball B branched along two arcs is a solid torus $\tilde{B}$, and a meridian for the solid torus is given by the preimage in $\tilde{B}$ of either of the arcs pushed out to the boundary of B . It follows that $\Sigma(K_-)$, $\Sigma(K_0)$, $\Sigma(K_+)$ each contain an embedded solid torus, such that the complements of these solid tori can be identified. The meridians which bound in $\Sigma(K_-)$, $\Sigma(K_0)$, $\Sigma(K_+)$ are shown in Figure 9. They may be oriented so that their homology classes intersect as follows:

$$(8) \quad \mu_- \cdot \mu_+ = 2, \quad \mu_+ \cdot \mu_0 = \mu_- \cdot \mu_0 = 1.$$

Suppose now that $K = K_-$ with $\sigma(K) = 2$, and K_+ is the unknot. Then $\Sigma(K_+) = S^3$, and $\Sigma(K_-)$ is $(2m - 1)/2$ surgery on some knot C . We wish to show that m is positive and even. For some longitude λ of C with $\mu_+ \cdot \lambda = 1$ we have

$$\mu_- = -2\lambda - (2m - 1)\mu_+.$$

Expressing λ in the basis μ_+ , μ_0 and plugging into (8) yields $\lambda = \mu_0 - m\mu_+$, from which we see that $\mu_0 = \lambda + m\mu_+$. In other words, $\Sigma(K_0)$ is m surgery on C .

We now use two properties of the Conway-normalised Alexander polynomial, c.f. [8]. Firstly, for a knot K , the sign of the Alexander polynomial at -1 is given by

$$(-1)^{\sigma(K)/2} \det K = \Delta_K(-1).$$




FIGURE 9. **Meridians in $\Sigma(K_-)$, $\Sigma(K_0)$, $\Sigma(K_+)$.** Rotating by π around the horizontal axis gives the solid torus as the double cover of a ball branched along the arcs in K_0 .

This shows that $\Delta_{K_+}(-1) = 1$ and $\Delta_{K_-}(-1) = -|2m - 1|$. Secondly we have the skein relation

$$\Delta_{K_+}(t) - \Delta_{K_-}(t) = (t^{-1/2} - t^{1/2})\Delta_{K_0}(t),$$

which yields

$$1 + |2m - 1| = 2|m|.$$

It follows that m and $2m - 1$ are both positive. Finally, the determinant and signature of a knot K are shown in [10, Theorem 5.6] to satisfy

$$(9) \quad \det(K) \equiv \sigma(K) + 1 \pmod{4},$$

from which it follows that $2m - 1$ is congruent to 3 modulo 4.

Similarly if K_- is the unknot and $\sigma(K_+) = 0$, we have that $\Sigma(K_+)$ is $(2m - 1)/2$ surgery on a knot C and we find $\Sigma(K_0)$ is $(m - 1)$ surgery on C . The skein relation gives $|2m - 1| - 1 = 2|m - 1|$, which again shows m is positive. From (9) we have $2m - 1$ is congruent to 1 modulo 4.

The general case follows easily from the above. Let $c_1, \dots, c_{p+n}$ be the set of crossings (p positive, n negative) in some chosen diagram that we change to unknot K . Then $\Sigma(K)$ is Dehn surgery on a link $L = C_1, \dots, C_{p+n}$, with linking matrix $\frac{1}{2}Q$. Each C_i corresponds to a crossing c_i . Dehn surgery on a sublink of L gives the double branched cover of a knot which is obtained from K by changing a subset of the crossings in $\mathcal{C}$. In particular $Q_{ii}/2$ surgery on the knot C_i yields the double branched cover of the knot K' which is obtained from K by changing all of the crossings except c_i . It follows from the unknotting number one case applied to K' that all diagonal entries of Q are positive and exactly those which correspond to negative crossings are congruent to 3 modulo 4.

It remains to prove that Q is positive-definite. Note that from (1) and the assumption $n = \sigma(K)/2$, the knot signature changes every time we change a negative crossing and is unchanged when we change a positive crossing. Let Q_k be the submatrix $(Q_{ij})_{i,j \leq k}$. Observe that since the off-diagonal entries are even, the determinant of Q_k is congruent modulo 4 to the product of the diagonal entries. Let K_k be the knot obtained from K by changing the crossings $c_{k+1}, \dots, c_{p+n}$. Suppose that $\det Q_{k-1}$ is positive, and hence equals $\det K_{k-1}$. If c_k is positive then

$$Q_{kk} \equiv 1 \implies \det Q_k \equiv \det Q_{k-1} \pmod{4}.$$

Also (9) implies that the determinants of K_k and K_{k-1} are congruent modulo 4. It follows that $\det Q_k \equiv \det K_k \pmod{4}$. Since $\det Q_k$ and $\det K_k$ are equal up to sign and odd, $\det Q_k$ must be positive.

On the other hand if c_k is a negative crossing then

$$\det Q_k \equiv \det Q_{k-1} + 2, \quad \det K_k \equiv \det K_{k-1} + 2 \pmod{4},$$

and we again find $\det Q_k$ to be positive. By induction $\det Q_k$ is positive for all k . $\square$

Finally note that we may reorient any of the link components $C_1, \dots, C_{p+n}$ without changing the resulting Dehn surgery. Also by rational handlesliding as in Proposition 2.1 we may change the linking matrix by “adding” $\pm 2C_j$ to C_i for any i, j . These operations preserve the congruence classes modulo 4 of the diagonal. The last claim in the statement of Theorem 3 now follows from the following lemma.

Lemma 3.3. *Any matrix $P \in GL(r, \mathbb{Z})$ which is congruent to the identity modulo 2 may be obtained from the identity by a sequence of row operations, each of which is either multiplying a row by -1 or adding an even multiple of one row to another.*

Proof. Let $\mathbf{b} = (b_1, \dots, b_r)$ be an element of $\mathbb{Z}^r$ with $\gcd(b_1, \dots, b_r) = 1$. Assume $b_i \geq 0$ for all i , and that b_1 is odd but the other components $b_2, \dots, b_r$ are even. Let b_j be the least positive component. By subtracting even multiples of b_j and then possibly changing sign, we may replace every other component b_i by b'_i , with $0 \leq b'_i \leq b_j$. By the gcd condition, the least positive b'_i is less than b_j unless $b_j = j = 1$. By iterating this procedure we see that $\mathbf{b}$ may be reduced to $(1, 0, \dots, 0)$.

Now suppose $P \in GL(r, \mathbb{Z})$ is congruent to I modulo 2, and let $\mathbf{b}$ be the first column of Q . The argument just given shows that Q may be replaced by a matrix with $(1, 0, \dots, 0)$ in the first column using the specified row operations. Then replacing the second column with $(*, 1, 0, \dots, 0)$ by row operations on the last $r - 1$ rows, and so on, we see that we may reduce P to I in this manner. $\square$

4. HEEGAARD FLOER HOMOLOGY

In this section we recall some properties of the Heegaard Floer homology invariants of Ozsváth and Szabó. Details are to be found in their papers, in particular [14, 15, 16].

Let Y be an oriented rational homology three-sphere. Recall that the space $\text{Spin}^c(Y)$ of spin^c structures on Y is isomorphic to $H^2(Y; \mathbb{Z})$. If $|H^2(Y; \mathbb{Z})|$ is odd then there is a canonical isomorphism which takes the unique spin structure to zero; this gives $\text{Spin}^c(Y)$ a group structure.

Fixing a spin^c structure $\mathfrak{s}$, the Heegaard Floer homology $HF^+(Y; \mathfrak{s})$ is a $\mathbb{Q}$ -graded abelian group with an action by $\mathbb{Z}[U]$, where U lowers the grading by 2. The *correction term* invariant is a rational number $d(Y, \mathfrak{s})$; it is defined to be the lowest grading of a nonzero homogeneous element of $HF^+(Y; \mathfrak{s})$ which is in the image of U^n for all $n \in \mathbb{N}$. These have the property that $d(Y, \mathfrak{s}) = -d(-Y, \mathfrak{s})$, where $-Y$ denotes Y with the opposite orientation. We will describe below how these correction terms may be computed in certain cases.

Now let X be a positive-definite four-manifold with boundary Y . Then it is shown in [14] that for any spin^c structure $\mathfrak{s}$ on X ,

$$(10) \quad c_1(\mathfrak{s})^2 - b_2(X) \geq 4d(Y, \mathfrak{s}|_Y),$$

$$(11) \quad \text{and } c_1(\mathfrak{s})^2 - b_2(X) \equiv 4d(Y, \mathfrak{s}|_Y) \pmod{2}.$$

This means that the correction terms of Y may be used to give an obstruction to Y bounding a four-manifold X with a given positive-definite intersection form. We will now elaborate on how this may be checked in practice.

Suppose for simplicity that X is simply-connected and that $|H^2(Y; \mathbb{Z})|$ is odd. Let r denote the second Betti number of X . Fix a basis for $H_2(X; \mathbb{Z})$ and thus an isomorphism

$$H_2(X; \mathbb{Z}) \cong \mathbb{Z}^r.$$

Let Q be the matrix of the intersection pairing of X in this basis; thus Q is a symmetric positive-definite $r \times r$ integer matrix with $\det Q = |H^2(Y; \mathbb{Z})|$. The dual basis gives an isomorphism between the second cohomology $H^2(X; \mathbb{Z})$ and $\mathbb{Z}^r$. The set $\{c_1(\mathfrak{s}) \mid \mathfrak{s} \in \text{Spin}^c(X)\} \subset H^2(X; \mathbb{Z})$ of first Chern classes of spin^c structures is equal to the set of characteristic covectors $\text{Char}(Q)$ for Q . These in turn are elements ξ of $\mathbb{Z}^r$ whose components ξ_i are congruent modulo 2 to the corresponding diagonal entries Q_{ii} of Q . The square of the first Chern class of a spin^c structure is computed using the pairing induced by Q on $H^2(X; \mathbb{Z})$; in our choice of basis this is given by $\xi^T Q^{-1} \xi$.

The long exact sequence of the pair (X, Y) yields the following short exact sequence:

$$0 \longrightarrow \mathbb{Z}^r \xrightarrow{Q} \mathbb{Z}^r \longrightarrow H^2(Y; \mathbb{Z}) \longrightarrow 0.$$

As in the introduction, define a function

$$m_Q : \mathbb{Z}^r / Q(\mathbb{Z}^r) \rightarrow \mathbb{Q}$$

by

$$m_Q(g) = \min \left\{ \frac{\xi^T Q^{-1} \xi - r}{4} \mid \xi \in \text{Char}(Q), [\xi] = g \right\}.$$

In computing m_Q it suffices to consider characteristic covectors $\xi = (\xi_1, \dots, \xi_r)$ whose components are smaller in absolute value than the corresponding diagonal entries of Q :

$$-Q_{ii} \leq \xi_i \leq Q_{ii}.$$

If, say, $\xi_i > Q_{ii}$, subtract twice the i th column of Q from ξ to see that $\xi^T Q^{-1} \xi$ is not minimal. A more difficult argument in [15] shows that it suffices to restrict to

$$-Q_{ii} \leq \xi_i \leq Q_{ii} - 2.$$

Thus it is straightforward, if tedious, to compute m_Q for a given positive-definite matrix Q .

The conditions (10) and (11) may now be expressed as follows:

Theorem 4.1 (Ozsváth-Szabó). *Let Y be a rational homology three-sphere which is the boundary of a simply-connected positive-definite four-manifold X , with $|H^2(Y; \mathbb{Z})|$ odd. If the intersection pairing of X is represented in a basis by the matrix Q then there exists a group isomorphism*

$$\phi : \mathbb{Z}^r / Q(\mathbb{Z}^r) \rightarrow \text{Spin}^c(Y)$$

with

$$(12) \quad m_Q(g) \geq d(Y, \phi(g)),$$

$$(13) \quad \text{and } m_Q(g) \equiv d(Y, \phi(g)) \pmod{2}$$

for all $g \in \mathbb{Z}^r / Q(\mathbb{Z}^r)$.

The four-manifold X is said to be *sharp* if equality holds in (12). In this case the correction terms for Y can be computed using the function m_Q described above. Also, if a rational homology sphere Y bounds a negative-definite four-manifold X such that $-X$ is sharp, then the correction terms for Y can be computed using the formula $d(Y, \mathfrak{s}) = -d(-Y, \mathfrak{s})$. Note that if K is a knot in S^3 then the standard orientation on S^3 induces an orientation on $\Sigma(K)$; letting $r(K)$ denote the reflection of K , we have $\Sigma(r(K)) \cong -\Sigma(K)$.

In particular let K be an alternating knot with double branched cover $\Sigma(K)$. Let G denote the positive-definite Goeritz matrix computed from an alternating diagram for K as follows. Colour the knot diagram in chessboard fashion according to the convention shown in Figure 10. (Note that this is the opposite convention to that used in [16], since they use the negative-definite Goeritz matrix.) Let $v_1, \dots, v_{k+1}$ denote the vertices of the white graph. Then G is the $k \times k$ symmetric matrix (g_{ij}) with entries

$$g_{ij} = \begin{cases} \text{the number of edges containing } v_i & \text{if } i = j \\ \text{minus the number of edges joining } v_i \text{ and } v_j & \text{if } i \neq j \end{cases}$$




FIGURE 10. Colouring convention for alternating knot diagrams.

for $i, j = 1, \dots, k$. It is shown in [16, Proposition 3.2] that G represents the intersection pairing of a sharp four-manifold bounded by $\Sigma(K)$. Thus the correction terms for $\Sigma(K)$ are given by m_G (for any choice of alternating diagram and any ordering of the white regions). Also it follows from [4] that with this colouring convention, the signature of K is given by

$$\sigma(K) = k - \mu,$$

where μ is the number of positive crossings in the alternating diagram used to compute G .

Also if K is a Montesinos knot then the double branched cover $\Sigma(K)$ is a Seifert fibred space which is given as the boundary of a plumbing of disk bundles over S^2 . This plumbing is determined (nonuniquely) by the Montesinos invariants which specify K . After possibly reflecting K we may choose the plumbing so that its intersection pairing is represented by a positive-definite matrix P . It is shown in [15] that the plumbing is sharp, so that the correction terms for $\Sigma(K)$ are given by m_P . (See [11] for a description of Montesinos knots and their branched double covers.)

Remark 4.2. *Checking the congruence condition (11) alone is equivalent to checking that the intersection pairing of X presents the linking pairing of Y ; see [13] for a detailed discussion.*

5. OBSTRUCTION TO UNKNOTTING

In this section we prove Theorems 1 and 5.

Let $\mathcal{Q}(r, \delta)$ denote the set of positive-definite symmetric integer matrices of rank r and determinant δ , on which $GL(r, \mathbb{Z})$ acts by $P \cdot Q = PQP^T$ with finite quotient (see e.g. [2]). Let $\mathcal{Q}(r, \delta)_2 \subset \mathcal{Q}(r, \delta)$ (resp. $GL(r, \mathbb{Z})_2 \subset GL(r, \mathbb{Z})$) denote the subset (resp. subgroup) consisting of matrices which are congruent to the identity modulo 2. Then the subset $\mathcal{Q}(r, \delta)_2/GL(r, \mathbb{Z})$ is clearly finite, and thus so is $\mathcal{Q}(r, \delta)_2/GL(r, \mathbb{Z})_2$ since $GL(r, \mathbb{Z})_2$ is a finite index subgroup of $GL(r, \mathbb{Z})$.

Proof of Theorem 5. By Theorem 3, the unknotting hypothesis implies that $\Sigma(K)$ is given by Dehn surgery on a link in S^3 with linking matrix $\frac{1}{2}Q_i$ for some i , where n of the diagonal entries of Q_i are congruent to 3 modulo 4. By Lemma 2.2, $\Sigma(K)$ bounds the 2-handlebody W specified by an integer-framed link with positive-definite linking matrix $\tilde{Q}_i$, which then represents the intersection pairing of W . The conclusion now follows from Theorem 4.1. $\square$

Proof of Theorem 1. Theorem 1 follows from Theorem 5 since a finite set of representatives of $\mathcal{Q}(2, \delta)_2/GL(2, \mathbb{Z})_2$ is given by the set of matrices

$$\left\{ Q = \begin{pmatrix} 2m_1 - 1 & 2a \\ 2a & 2m_2 - 1 \end{pmatrix} \mid \det Q = \delta, 0 \leq a < m_1 \leq m_2 \right\},$$

and since the correction terms $d(\Sigma(K), \mathfrak{s})$ may be computed using a positive-definite Goeritz matrix G when K is alternating. $\square$

Remark 5.1. *Theorems 1 and 5 do not use all of the information from Theorem 3. We have only used the information about the intersection pairing of the four-manifold W bounded by $\Sigma(K)$, and not the fact that W is a surgery cobordism arising from a half-integral surgery. Comparing to Theorem 1.1 in [16], we have generalised conditions (1) and (2) to the case of $u(K) > 1$ but not the symmetry condition (3). It is to be hoped that the symmetry condition may also be generalised in some way, leading to a stronger obstruction and computation of some more unknotting numbers.*

6. EXAMPLES

Proof of Corollary 2. For each knot in Corollary 2 we distinguish between K and its reflection $r(K)$ by specifying that K has positive signature.

We start with the knot $K = 9_{10}$ shown in Figure 11. This is the two-bridge knot $S(33, 23)$. It has signature 4, and it is easy to see that 3 crossing changes suffice to unknot it. Thus the unknotting number is either 2 or 3, and if it can be unknotted by changing two crossings then both are negative ($p = 0$ and $n = 2$).

With the white regions labelled as shown in the figure, the Goeritz matrix is

$$G = \begin{pmatrix} 4 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 4 \end{pmatrix}.$$

Using m_G , we find the correction terms of $\Sigma(K)$ to be:

$$A = \left\{ \begin{array}{cccccccccccc} -1, & -\frac{23}{33}, & \frac{7}{33}, & -\frac{3}{11}, & -\frac{5}{33}, & \frac{19}{33}, & -\frac{1}{11}, & -\frac{5}{33}, & \frac{13}{33}, & -\frac{5}{11}, & -\frac{23}{33}, \\ -\frac{1}{3}, & \frac{7}{11}, & \frac{7}{33}, & \frac{13}{33}, & \frac{13}{11}, & \frac{19}{33}, & \frac{19}{33}, & \frac{13}{11}, & \frac{13}{33}, & \frac{7}{33}, & \frac{7}{11}, \\ -\frac{1}{3}, & -\frac{23}{33}, & -\frac{5}{11}, & \frac{13}{33}, & -\frac{5}{33}, & -\frac{1}{11}, & \frac{19}{33}, & -\frac{5}{33}, & -\frac{3}{11}, & \frac{7}{33}, & -\frac{23}{33} \end{array} \right\}.$$

The order of this list corresponds to the cyclic group structure of $\text{Spin}^c(\Sigma(K)) \cong H^2(\Sigma(K); \mathbb{Z})$, and the first element is the correction term of the spin structure.




FIGURE 11. **The knot** $9_{10} = S(33, 23)$. Note that changing the circled crossings will give the unknot. The labels $v_1, \dots, v_5$ correspond to vertices of the white graph.

The determinant of 9_{10} is 33. To find a matrix $\tilde{Q}$ as in Theorem 1 we need to find (m_1, a, m_2) with

$$(2m_1 - 1)(2m_2 - 1) - 4a^2 = 33,$$

$$0 \leq a < m_1 \leq m_2,$$

and m_1 and m_2 are even. There are two solutions: $(2, 0, 6)$ and $(4, 2, 4)$. Computing $m_{\tilde{Q}}$ for each of the matrices

$$\tilde{Q}_1 = \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 0 & 0 & 6 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix}, \quad \tilde{Q}_2 = \begin{pmatrix} 4 & 1 & 2 & 0 \\ 1 & 2 & 0 & 0 \\ 2 & 0 & 4 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

yields the following lists:

$$B_1 = \left\{ \begin{array}{cccccccccccc} -1, & -\frac{5}{33}, & \frac{13}{33}, & \frac{7}{11}, & \frac{19}{33}, & \frac{7}{33}, & -\frac{5}{11}, & \frac{19}{33}, & \frac{43}{33}, & -\frac{3}{11}, & -\frac{5}{33}, \\ -\frac{1}{3}, & -\frac{9}{11}, & \frac{13}{33}, & \frac{43}{33}, & -\frac{1}{11}, & \frac{7}{33}, & \frac{7}{33}, & -\frac{1}{11}, & \frac{43}{33}, & \frac{13}{33}, & -\frac{9}{11}, \\ -\frac{1}{3}, & -\frac{5}{33}, & -\frac{3}{11}, & \frac{43}{33}, & \frac{19}{33}, & -\frac{5}{11}, & \frac{7}{33}, & \frac{19}{33}, & \frac{7}{11}, & \frac{13}{33}, & -\frac{5}{33} \end{array} \right\},$$

$$B_2 = \left\{ \begin{array}{cccccccccccc} -1, & -\frac{19}{33}, & \frac{23}{33}, & \frac{9}{11}, & -\frac{7}{33}, & -\frac{13}{33}, & \frac{3}{11}, & -\frac{7}{33}, & \frac{5}{33}, & -\frac{7}{11}, & -\frac{19}{33}, \\ \frac{1}{3}, & \frac{1}{11}, & \frac{23}{33}, & \frac{5}{33}, & \frac{5}{11}, & -\frac{13}{33}, & -\frac{13}{33}, & \frac{5}{11}, & \frac{5}{33}, & \frac{23}{33}, & \frac{1}{11}, \\ \frac{1}{3}, & -\frac{19}{33}, & -\frac{7}{11}, & \frac{5}{33}, & -\frac{7}{33}, & \frac{3}{11}, & -\frac{13}{33}, & -\frac{7}{33}, & \frac{9}{11}, & \frac{23}{33}, & -\frac{19}{33} \end{array} \right\}.$$

We claim that for both $\tilde{Q}_1$ and $\tilde{Q}_2$ it is impossible to find a group automorphism ϕ of $\mathbb{Z}/33$ satisfying the required inequality and congruence conditions. This is immediate in either case by considering the minimal elements (excluding -1 which appears in all 3 lists). We have the entry $-9/11$ in B_1 . By inspection there is no element in A which is less than or equal to $-9/11$, and differs from it by a multiple of 2. The same applies to $-7/11$ in B_2 . We conclude that 9_{10} cannot be unknotted by two crossing changes and $u(9_{10}) = 3$.

Similar calculations show that $9_{13}, 9_{38}, 10_{53}, 10_{101}$ and 10_{120} cannot be unknotted with two crossing changes. All of these knots are alternating, have signature four and cyclic $H^2(\Sigma(K); \mathbb{Z})$. By inspection of their diagrams (see e.g. [1]), all can be unknotted with three crossing changes. For some details of the calculations for these knots, see Table 1. Note that we use the knot diagrams from [1] to compute the Goeritz matrices for these knots, after possibly reflecting to ensure positive signature.

Finally consider $K = 9_{35}$, pictured in Figure 12. It has signature 2 and can be unknotted with 3 crossing changes. The Goeritz matrix from the figure is

$$G = \begin{pmatrix} 6 & -3 \\ -3 & 6 \end{pmatrix}.$$

We note that this presents $H^2(\Sigma(K); \mathbb{Z})$ which is thus 2-cyclic; this shows (by Montesinos' theorem for example but by an inequality originally due to Wendt) that $u(K) \geq 2$. We can use m_G to compute the correction terms of $\Sigma(K)$, which are

$$A = \begin{bmatrix} -\frac{1}{2} & \frac{19}{18} & -\frac{5}{18} & \frac{3}{2} & \frac{7}{18} & \frac{7}{18} & \frac{3}{2} & -\frac{5}{18} & \frac{19}{18} \\ \frac{1}{6} & -\frac{5}{18} & \frac{7}{18} & \frac{1}{6} & \frac{19}{18} & \frac{19}{18} & \frac{1}{6} & \frac{7}{18} & -\frac{5}{18} \\ \frac{1}{6} & -\frac{5}{18} & \frac{7}{18} & \frac{1}{6} & \frac{19}{18} & \frac{19}{18} & \frac{1}{6} & \frac{7}{18} & -\frac{5}{18} \end{bmatrix}.$$

Here the rectangular array shows the $\mathbb{Z}/3 \oplus \mathbb{Z}/9$ group structure; the top left entry is the correction term of the spin structure.




FIGURE 12. **The Montesinos knot** $9_{35} = M(0; (3, 1), (3, 1), (3, 1))$.


Suppose that 9_{35} may be unknotted by changing one positive and one negative crossing. The only matrix which satisfies the conditions of Theorem 1 and which presents $\mathbb{Z}/3 \oplus \mathbb{Z}/9$ is

$$\tilde{Q} = \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 0 & 0 & 5 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix}.$$

Computing $m_{\tilde{Q}}$ yields another array whose minimal entry is $-17/18$; we conclude that there is no automorphism ϕ of $\mathbb{Z}/3 \oplus \mathbb{Z}/9$ satisfying the conclusion of Theorem 1.

This is not enough to rule out the possibility that $u(9_{35}) = 2$; it does however show that if 9_{35} can be unknotted by two crossing changes, then they are both negative crossings. Using the value of the Jones polynomial at $e^{i\pi/3}$, Traczyk has shown in [18] that if 9_{35} can be unknotted by changing two crossings, then the crossings have different signs. We conclude that $u(9_{35}) = 3$. $\square$

Proof of Corollary 6. The two-bridge knot $K = S(51, 35)$ is listed in [1] as 11a365 and is shown in Figure 13. It has signature 6, and from the diagram we see that it may be unknotted by changing 4 crossings. We will apply Theorem 5 to show that it does not have $u(K) = n = 3$.

FIGURE 13. **The two-bridge knot $S(51, 35)$, or $11a365$.**

Note that $\det K = 51$. In order to apply Theorem 5 we first need to find a set of representatives of the finite quotient $\mathcal{Q}(3, 51)_2/GL(3, \mathbb{Z})_2$ with all diagonal entries conjugate to 3 modulo 4. According to [6], a complete set of representatives of $\mathcal{Q}(3, 51)/GL(3, \mathbb{Z})$ is given by the (Eisenstein reduced) matrices

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 51 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 1 & 26 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 17 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 4 & 1 \\ 0 & 1 & 13 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 5 & 2 \\ 0 & 2 & 11 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 6 & 3 \\ 0 & 3 & 10 \end{pmatrix},$$

$$\begin{pmatrix} 2 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 9 \end{pmatrix}, \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 17 \end{pmatrix}, \begin{pmatrix} 3 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 6 \end{pmatrix}, \begin{pmatrix} 3 & 1 & 1 \\ 1 & 4 & 0 \\ 1 & 0 & 5 \end{pmatrix}, \begin{pmatrix} 4 & 1 & 2 \\ 1 & 4 & 2 \\ 2 & 2 & 5 \end{pmatrix}.$$

Note that if $P \in GL(3, \mathbb{Z})$ satisfies $PP^T \equiv I \pmod{2}$, then P is conjugate to a permutation matrix modulo 2. Thus if $P \in GL(3, \mathbb{Z})$ and $Q, PQP^T \in \mathcal{Q}(3, 51)_2$ then Q and PQP^T have the same number of diagonal entries conjugate to 3 modulo 4. We

therefore eliminate the forms represented by $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 51 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 17 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 5 & 2 \\ 0 & 2 & 11 \end{pmatrix}.$

For each remaining form in the list, we look for a basis in which the form is congruent to the identity modulo 2. If no such basis exists, or if we find that some diagonal entry is not conjugate to 3 modulo 4, we eliminate the form. This leaves us with

the following four forms to consider: $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 1 & 26 \end{pmatrix} \sim \begin{pmatrix} 3 & 2 & 0 \\ 2 & 27 & 26 \\ 0 & 26 & 27 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 6 & 3 \\ 0 & 3 & 10 \end{pmatrix} \sim$
 $\begin{pmatrix} 11 & 4 & -6 \\ 4 & 7 & 4 \\ -6 & 4 & 11 \end{pmatrix}, \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 17 \end{pmatrix} \sim \begin{pmatrix} 19 & 18 & 18 \\ 18 & 19 & 16 \\ 18 & 16 & 19 \end{pmatrix}, \text{ and } \begin{pmatrix} 3 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 6 \end{pmatrix} \sim \begin{pmatrix} 3 & 0 & -2 \\ 0 & 3 & 0 \\ -2 & 0 & 7 \end{pmatrix}.$

From Figure 13 we may write down the Goeritz matrix G ; the correction terms $\{d(\Sigma(K), \mathfrak{s})\}$ are then given by m_G . For each Q in

$$\left\{ \begin{pmatrix} 3 & 2 & 0 \\ 2 & 27 & 26 \\ 0 & 26 & 27 \end{pmatrix}, \begin{pmatrix} 11 & 4 & -6 \\ 4 & 7 & 4 \\ -6 & 4 & 11 \end{pmatrix}, \begin{pmatrix} 19 & 18 & 18 \\ 18 & 19 & 16 \\ 18 & 16 & 19 \end{pmatrix}, \begin{pmatrix} 3 & 0 & -2 \\ 0 & 3 & 0 \\ -2 & 0 & 7 \end{pmatrix} \right\},$$

one may check that there is no isomorphism

$$\phi : \Gamma_{\tilde{Q}} \rightarrow \text{Spin}^c(\Sigma(K))$$

satisfying the conclusion of Theorem 5. We conclude that the unknotting number of K is 4. $\square$

Remark 6.1. *In the last step of the proof of Corollary 6 it is much quicker in some cases to change basis before computing $m_{\tilde{Q}}$, so as to work with a matrix with smaller*

diagonal entries. For example with $Q = \begin{pmatrix} 19 & 18 & 18 \\ 18 & 19 & 16 \\ 18 & 16 & 19 \end{pmatrix}$, we have

$$\tilde{Q} = \begin{pmatrix} 10 & 1 & 9 & 0 & 9 & 0 \\ 1 & 2 & 0 & 0 & 0 & 0 \\ 9 & 0 & 10 & 1 & 8 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 \\ 9 & 0 & 8 & 0 & 10 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{pmatrix} \sim \begin{pmatrix} 10 & 1 & -1 & 0 & -1 & 0 \\ 1 & 2 & -1 & 0 & -1 & 0 \\ -1 & -1 & 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 \\ -1 & -1 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{pmatrix},$$

by subtracting the first basis vector from the third and fifth. As a result we need to consider $2^5 \cdot 10$ characteristic covectors to compute $m_{\tilde{Q}}$ instead of $2^3 \cdot 10^3$.

Knot	Goeritz matrix	$\min_{g \neq 0} \{m_G(g)\}$	(m_1, a, m_2)	$\min_{g \neq 0} \{m_Q(g)\}$
9_{13}	$\begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 4 & -1 \\ 0 & 0 & -1 & 4 \end{pmatrix}$	$-\frac{27}{37}$	$(10, 9, 10)$	$-\frac{33}{37}$
9_{38}	$\begin{pmatrix} 4 & -1 & -1 & 0 \\ -1 & 4 & -2 & 0 \\ -1 & -2 & 4 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$	$-\frac{37}{57}$	$(2, 0, 10)$	$-\frac{51}{57}$
			$(6, 4, 6)$	$-\frac{45}{57}$
10_{53}	$\begin{pmatrix} 4 & -1 & 0 & 0 \\ -1 & 4 & -1 & -1 \\ 0 & -1 & 4 & -1 \\ 0 & -1 & -1 & 2 \end{pmatrix}$	$-\frac{53}{73}$	$(4, 1, 6)$	$-\frac{59}{73}$
10_{101}	$\begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 4 & -1 & -1 \\ 0 & -1 & 4 & -1 \\ 0 & -1 & -1 & 4 \end{pmatrix}$	$-\frac{59}{85}$	$(6, 3, 6)$	$-\frac{65}{85}$
			$(22, 21, 22)$	$-\frac{81}{85}$
10_{120}	$\begin{pmatrix} 4 & -2 & 0 & -1 \\ -2 & 4 & -1 & 0 \\ 0 & -1 & 4 & -2 \\ -1 & 0 & -2 & 4 \end{pmatrix}$	$-\frac{69}{105}$	$(2, 0, 18)$	$-\frac{99}{105}$
			$(4, 0, 8)$	$-\frac{91}{105}$
			$(6, 2, 6)$	$-\frac{83}{105}$
			$(10, 8, 10)$	$-\frac{93}{105}$

TABLE 1. **Data for knots in Corollary 2.** The fourth column contains possible coefficients of the matrix Q in Theorem 1.

REFERENCES

- [1] J. C. Cha & C. Livingston, *Table of knot invariants*, <http://www.indiana.edu/~knotinfo>.
- [2] J. W. S. Cassels, *Rational quadratic forms*, Academic Press, 1978.
- [3] T. D. Cochran & W. B. R. Lickorish, *Unknotting information from 4-manifolds*, Trans. Amer. Math. Soc. **297** 1986, 125–142.
- [4] C. McA. Gordon & R. A. Litherland, *On the signature of a link*, Invent. Math. **47** 1978, 53–69.
- [5] R. E. Gompf & A. I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Math. **20**, Amer. Math. Soc., 1999.
- [6] B. W. Jones, *A table of Eisenstein-reduced positive ternary quadratic forms of determinant ≤ 200* , National Research Council Bulletin **97** 1935.
- [7] W. B. R. Lickorish, *The unknotting number of a classical knot*, Contemp. Math. **44** 1985, 117–121.
- [8] W. B. R. Lickorish, *An introduction to knot theory*, Graduate Texts in Math. **175**, Springer, 1997.
- [9] J. M. Montesinos, *Surgery on links and double branched coverings of S^3* , Ann. of Math. Studies **84** 1975, 227–259.
- [10] K. Murasugi, *On a certain numerical invariant of link types*, Trans. Amer. Math. Soc. **117** 1965, 387–422.
- [11] B. Owens & S. Strle. *Rational homology spheres and the four-ball genus of knots*, math.GT/0308073, to appear in Advances in Mathematics.
- [12] B. Owens & S. Strle. *A characterisation of the $n\langle 1 \rangle \oplus \langle 3 \rangle$ form and applications to rational homology spheres*, math.GT/0312266.
- [13] B. Owens & S. Strle. *Definite manifolds bounded by rational homology three spheres*, to appear in the Proceedings of the Fields-McMaster Conference on Geometry and Topology of Manifolds.
- [14] P. Ozsváth & Z. Szabó. *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Advances in Mathematics **173** 2003, 179–261.
- [15] P. Ozsváth & Z. Szabó. *On the Floer homology of plumbed three-manifolds*, Geometry and Topology **7** 2003, 225–254.
- [16] P. Ozsváth & Z. Szabó. *Knots with unknotting number one and Heegaard Floer homology*, math.GT/0401426.
- [17] A. Stoimenow. *Polynomial values, the linking form and unknotting numbers*, Math. Res. Lett. **11** 2004, 755–769.
- [18] P. Traczyk. *A criterion for signed unknotting number*, Low-dimensional topology (Funchal, 1998), Contemp. Math. **233**, Amer. Math. Soc., 1999, 215–220.